

# Scalar dark matter

work with

Bohdan Grzadkowski, Odd Magne Ogreid,  
Alexander Pukhov, Madhi Purmohammadi

Per Osland

August 2011

Warszawa

PRD 2009, JHEP 2011



# Dark Matter

- Astrophysical evidence: 85% of Universe matter is dark [list from Hooper & Baltz, 2008]
  - rotational speeds of galaxies
  - orbital velocities of galaxies within clusters
  - gravitational lensing
  - cosmic microwave background
  - light element abundance
  - large scale structure
- Not homogeneously distributed
- Many particle and astrophysical candidates



# Dark Matter

- Astrophysical evidence: 85% of Universe matter is dark [list from Hooper & Baltz, 2008]
  - rotational speeds of galaxies
  - orbital velocities of galaxies within clusters
  - gravitational lensing
  - cosmic microwave background
  - light element abundance
  - large scale structure
- Not homogeneously distributed
- Many particle and astrophysical candidates

Local halo density:  $0.22\text{--}0.75 \text{ GeV/cm}^3$



# Dark Matter

- Astrophysical evidence: 85% of Universe matter is dark [list from Hooper & Baltz, 2008]
  - rotational speeds of galaxies
  - orbital velocities of galaxies within clusters
  - gravitational lensing
  - cosmic microwave background
  - light element abundance
  - large scale structure
- Not homogeneously distributed
- Many particle and astrophysical candidates

Local halo density:  $0.22\text{--}0.75 \text{ GeV/cm}^3$       One per cup of coffee

# footnote: scales

$$1 \text{ light year} \approx (3 \times 10^8 \text{ m/s}) \times (\pi \times 10^7 \text{ s}) \approx 10^{16} \text{ m}$$

$$\text{galactic diameter} \approx 80,000 \text{ light years} \approx 10^{21} \text{ m}$$

$$1 \text{ parsec} = 1 \text{ pc} \approx 3.26 \text{ light years}$$

Solar system:  $\sim 8.5 \text{ kpc}$  from Galactic Center

$8.5 \text{ kpc}$  ( $\sim 28,000 \text{ light years}$ ) vs  $40,000 \text{ light years}$

➡ We are far from the galactic center

# Scales

Unknown

Planck scale



Particle physics

Other interactions explored down to here

Gravity explored down to here

Atomic physics

You are here

Solar system

light year

Galaxies  
Clusters of galaxies

$10^{-36}$  m


$10^{-24}$  m

$10^{-12}$  m

1 m


$10^{12}$  m

$10^{24}$  m




# Ambition:

- Problem at scales of  $10^{20}$  m
- Solution: new physics at scales below  $10^{-18}$  m




# Schematic spectrum





# The Big Picture



- neutrino  $\nu$  – hot DM
- neutralino  $\chi$
- “generic” WIMP
- axion  $a$
- axino  $\tilde{a}$
- gravitino  $\tilde{G}$
- ????

solution of DM: must go beyond SM!

Leszek Roszkowski, 2010

# Popular Dark Matter candidates

- Neutralino of Supersymmetry

- Axions, axinos

- Neutrinos (exist!)

- Gravitinos

- Scalars

# Simple estimate

Density of dark matter given by Early Universe consideration:  
Equilibrium between Hubble expansion and annihilation

Calculable

$$\Omega_{\text{DM}} h^2 = C \frac{T_0^3}{M_{\text{Pl}}^3} \frac{1}{\langle \sigma v \rangle}$$

$T_0$  CMBR temp

WMAP:  $\Omega_{\text{DM}} h^2 = 0.1131 \pm 0.0034$  and  $\langle \sigma v \rangle = \frac{\alpha^2}{m^2} \Rightarrow m \sim 100 \text{ GeV}$

mass scale fits well with SUSY



# Direct/indirect detection?

- Interacts very weakly with ordinary matter
- No strong (nuclear) interactions
- Might be local enhancement near solar system
- Ordinary particles might be scattered, recoil (CDMS-II)
- Might annihilate in sun, look for photons/positrons
- Might annihilate in space, Milky Way,...

PAMELA, ATIC, Fermi-LAT,...

# Scalar DM

- “Inert (Scalar) Doublet Model”, Barbieri et al, 2006

Extend SM with additional scalar doublet, unbroken  $Z_2$  symmetry makes lightest “odd” particle stable. No vev, no direct coupling to SM matter.

$$V = \mu_1^2 |H_1|^2 + \mu_2^2 |H_2|^2 + \lambda_1 |H_1|^4 + \lambda_2 |H_2|^4 \\ + \lambda_3 |H_1|^2 |H_2|^2 + \lambda_4 |H_1^\dagger H_2|^2 + \frac{\lambda_5}{2} \left[ (H_1^\dagger H_2)^2 + h.c. \right]$$

## 2HDM

Unbroken  $Z_2$ :  $H_1 \rightarrow H_1$  and  $H_2 \rightarrow -H_2$ . Coupling to non-inert Higgs  
 $\langle H_2 \rangle = 0$ .  $\lambda_L = (\lambda_3 + \lambda_4 + \lambda_5)/2$

# Motivation (Barbieri et al)

May alleviate Little Hierarchy Problem,  
by allowing heavier SM Higgs (400 GeV)  
without conflict with “electroweak  
precision data” (S and T).

Also work by:

- Ma; Kubo, Ma, Suematso; Cao, Ma, Rajasekaran
- Lopez Honorez, Nezri, Oliver, Tytgat
- Gustafsson, Lundstrom, Bergstrom, Edsjo
- Cirelli, Strumia, Tamburini
- Andreas, Hambye, Tytgat
- Dolle, Su
- Pierce, Thaler
- ...



# Motivation (Barbieri et al)

May alleviate Little Hierarchy Problem,  
by allowing heavier SM Higgs (400 GeV)  
without conflict with “electroweak  
precision data” (S and T).

New particles (from isospin doublet):

$H^\pm$

$A_0$

$H_0$  Stable - DM

$A_0$


$H^\pm$

$H_0$  Stable - DM


# Motivation (Barbieri et al)

May alleviate Little Hierarchy Problem,  
by allowing heavier SM Higgs (400 GeV)  
without conflict with “electroweak  
precision data” (S and T).

New particles (from isospin doublet):



Stable - DM



Stable - DM

Laura Lopez Honorez, Emmanuel Nezri, Josep F. Oliver, Michel H.G. Tytgat




Figure 1: Annihilation channels into gauge bosons final state with corresponding couplings.




Figure 2: Annihilation channels into Higgs final state.




# Allowed regions

$$M_{H^\pm} > 79.3 \text{ GeV}$$

LEP (PDG)

$$M_A \gtrsim 110 \text{ GeV}$$

LEP, Lundquist et al



Three (4) regions allowed for  $M_S \equiv M_{H_0}$

LOW: Andreas, Hambye, Tytgat; Hambye, Tytgat

MEDIUM: Barbieri, Hall, Rychkov; Lopez Honorez, Nezri, Oliver, Tytgat

“NEW, VIABLE”: Lopez Honorez, Yaguna

HIGH: Lopez Honorez, Nezri, Oliver, Tytgat; Hambye, Ling, Lopez Honorez, Rocher; Cirelli, Fornengo, Strumia

# IDM2: 2HDM + inert doublet

Grzadkowski et al, 2009

## Motivation: IDM + CP violation

Fields:

$$\Phi_1 = \begin{pmatrix} \varphi_1^+ \\ (v_1 + \eta_1 + i\chi_1)/\sqrt{2} \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} \varphi_2^+ \\ (v_2 + \eta_2 + i\chi_2)/\sqrt{2} \end{pmatrix}$$

$$\eta = \begin{pmatrix} \eta^+ \\ (S + iA)/\sqrt{2} \end{pmatrix}$$

Potential:

$$V(\Phi_1, \Phi_2, \eta) = V_{12}(\Phi_1, \Phi_2) + V_3(\eta) + V_{123}(\Phi_1, \Phi_2, \eta)$$

Coupling:



$$V_{12}(\Phi_1, \Phi_2) = -\frac{1}{2} \left\{ m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 + \left[ m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.} \right] \right\}$$

(standard)

$$+ \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) \\ + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \frac{1}{2} \left[ \lambda_5 (\Phi_1^\dagger \Phi_2)^2 + \text{h.c.} \right]$$

$$V_3(\eta) = m_\eta^2 \eta^\dagger \eta + \frac{\lambda_\eta}{2} (\eta^\dagger \eta)^2$$

**Coupling:**

(most general)

$$V_{123}(\Phi_1, \Phi_2, \eta) = \lambda_{1133} (\Phi_1^\dagger \Phi_1) (\eta^\dagger \eta) + \lambda_{2233} (\Phi_2^\dagger \Phi_2) (\eta^\dagger \eta) \\ + \lambda_{1331} (\Phi_1^\dagger \eta) (\eta^\dagger \Phi_1) + \lambda_{2332} (\Phi_2^\dagger \eta) (\eta^\dagger \Phi_2) \\ + \frac{1}{2} \left[ \lambda_{1313} (\Phi_1^\dagger \eta)^2 + \text{h.c.} \right] + \frac{1}{2} \left[ \lambda_{2323} (\Phi_2^\dagger \eta)^2 + \text{h.c.} \right]$$

**Many parameters...**



## Many parameters! Simplify!

“Dark democracy”:  $\lambda_a \equiv \lambda_{1133} = \lambda_{2233},$   
 $\lambda_b \equiv \lambda_{1331} = \lambda_{2332},$   
 $\lambda_c \equiv \lambda_{1313} = \lambda_{2323} \text{ (real)}.$

Masses of inert sector:

$$\begin{aligned} M_{\eta^\pm}^2 &= m_\eta^2 + \frac{1}{2}\lambda_a v^2, \\ M_S^2 &= m_\eta^2 + \frac{1}{2}(\lambda_a + \lambda_b + \lambda_c)v^2 = M_{\eta^\pm}^2 + \frac{1}{2}(\lambda_b + \lambda_c)v^2, \\ M_A^2 &= m_\eta^2 + \frac{1}{2}(\lambda_a + \lambda_b - \lambda_c)v^2 = M_{\eta^\pm}^2 + \frac{1}{2}(\lambda_b - \lambda_c)v^2 \end{aligned}$$


**Important:**

These  $\lambda_{a,b,c}$  characterize **coupling** of inert sector to non-inert sector, and also **mass splitting** in inert sector

# “Dark democracy”

## DOCTOR FUN

20 June 2003



Copyright © 2003 David Farley, d-farley@ibiblio.org  
<http://ibiblio.org/Dave/drfun.html>

This cartoon is made available on the Internet for personal viewing only. Opinions expressed herein are solely those of the author.

"If we held free elections they'd just elect a cat."

# Constraints

- positivity (rather complicated), 20% excluded
- unitarity, 60% excluded
- global minimum, 10% excluded
- additional 2HDM constraints:  $T$ ,  $b \rightarrow s\gamma$  etc

• DM  
determined by MicrOMEGAs

EW “precision data”

# Positivity

Define:

$$\lambda_x = \lambda_3 + \min(0, \lambda_4 - |\lambda_5|)$$

$$\lambda_y = \lambda_{1133} + \min(0, \lambda_{1331} - |\lambda_{1313}|)$$

$$\lambda_z = \lambda_{2233} + \min(0, \lambda_{2332} - |\lambda_{2323}|)$$

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_\eta > 0, \quad \lambda_x > -\sqrt{\lambda_1 \lambda_2}$$

$$\lambda_y > -\sqrt{\lambda_1 \lambda_\eta}, \quad \lambda_z > -\sqrt{\lambda_2 \lambda_\eta}$$

Plus additional constraint, which in the case of  
Dark democracy  $\lambda_y = \lambda_z$  takes the form:

$$\lambda_y \geq 0 \vee \left( \lambda_\eta \lambda_x - \lambda_y^2 > -\sqrt{(\lambda_\eta \lambda_1 - \lambda_y^2)(\lambda_\eta \lambda_2 - \lambda_y^2)} \right)$$

# Getting correct DM density

Main Early Universe annihilation mechanisms:

- Annihilation to  $W^+ W^-$ , effective above 75 GeV
- Annihilation via real or virtual neutral Higgs

like IDM...



# Annihilation in the Early Universe

The DM particles can annihilate via the gauge coupling:

$$SSW^+W^- : \frac{ig^2}{2}$$
$$SSZZ : \frac{ig^2}{2\cos^2\theta_W}$$

# Annihilation in the Early Universe

The DM particles can annihilate via the gauge coupling:

$$SSW^+W^- : \quad \frac{ig^2}{2}$$

$$SSZZ : \quad \frac{ig^2}{2\cos^2\theta_W}$$

or to non-inert scalars via trilinear

$$SSH_j : \quad -2iF_{SSj}\lambda_L v$$

**Note**



$$F_{SSj} = \cos\beta R_{j1} + \sin\beta R_{j2}$$



or quartic couplings:

$$SSH_j H_j : \quad -2i(\lambda_L - \lambda_c R_{j3}^2)$$

$$SSH_j H_k : \quad 2i\lambda_c R_{j3} R_{k3}$$

$$SSH^+ H^- : \quad -i\lambda_a$$

# Allowed regions in $M_s$




# Scan over parameters

1.  $M_S, M_1$  (lowest masses of inert and 2HDM sectors, fixed)
2.  $M_A, M_{\eta^\pm}$  (inert sector, physical masses, fixed).
3.  $M_2, \mu$  (2HDM sector parameters)
4.  $m_\eta$  (inert sector, soft mass parameter, fixed).
5.  $\tan \beta, M_{H^\pm}$  (2HDM sector),  
 $0.5 \leq \tan \beta \leq 50, 300 \text{ GeV} \leq M_{H^\pm} \leq 700 \text{ GeV}.$
6.  $\alpha_1, \alpha_2, \alpha_3$  (2HDM sector),  
 $-\pi/2 \leq \alpha_{1,2} \leq \pi/2$ , and  $0 \leq \alpha_3 \leq \pi/2.$

**Collect results in  $M_{\eta^\pm}, m_\eta$  plane**


Light DM: coupling to Higgs,  $|\lambda_L|$  can not be too small






$2 \times 60 \text{ GeV} = 120 \text{ GeV}$  resonant annihilation


$\lambda_L$  small



Mass of charged state





forbidden,  
coannihilation too fast



Coannihilation just right:


$$S\eta^\pm \rightarrow W^\pm \gamma$$

$$S\eta^\pm \rightarrow W^\pm{}^* \rightarrow u\bar{d}, d\bar{u}, c\bar{s}, s\bar{c}$$




Mass of charged state

# Allowed regions in $M_S, M_1$





# Allowed regions in $M_S, M_1$







# High region:



# High region:



Note small mass differences

# CP violation

Measured in terms of invariants (Gunion and Haber, 2005):

$$\text{Im } J_1 = -\frac{v_1^2 v_2^2}{v^4} (\lambda_1 - \lambda_2) \text{Im } \lambda_5$$

$$\text{Im } J_2 = -\frac{v_1^2 v_2^2}{v^8} \left[ \left( (\lambda_1 - \lambda_3 - \lambda_4)^2 - |\lambda_5|^2 \right) v_1^4 + 2(\lambda_1 - \lambda_2) \text{Re } \lambda_5 v_1^2 v_2^2 \right. \\ \left. - \left( (\lambda_2 - \lambda_3 - \lambda_4)^2 - |\lambda_5|^2 \right) v_2^4 \right] \text{Im } \lambda_5$$

$$\text{Im } J_3 = \frac{v_1^2 v_2^2}{v^4} (\lambda_1 - \lambda_2) (\lambda_1 + \lambda_2 + 2\lambda_4 + 2\lambda_b) \text{Im } \lambda_5$$

 small extra contribution


CP violation if at least one of these is non-zero

# CP violation

A “representative” case:

$$(M_S, M_A, M_{\eta^\pm}, m_\eta) = (75, 110, 90, 100) \text{ GeV}$$

$$(M_1, M_2, \mu) = (120, 300, 200) \text{ GeV}$$




orders of magnitude higher than in SM


(No solution found for large  $\tan\beta$ )

# CP violation: electron edm

Unit:  $e \cdot 10^{-27} \text{ cm}$  ( $\sim$  current limit)




# Direct detection







# Direct detection



# Compare CDMS-II and XENON100



# Compare CDMS-II and XENON100



# LHC prospects

If charged and neutral scalars of inert doublet are at electroweak scale, then scalars can be produced and perhaps even observed at the LHC:

$$pp \rightarrow SSX, AAX, SAX, S\eta^\pm X, A\eta^\pm X, \eta^+ \eta^- X.$$

# LHC prospects

If charged and neutral scalars of inert doublet are at electroweak scale, then scalars can be produced and perhaps even observed at the LHC:

$$pp \rightarrow SSX, AAX, SAX, S\eta^\pm X, A\eta^\pm X, \eta^+ \eta^- X.$$



followed by:

$$\eta^+ \rightarrow S\ell^+ \nu_\ell$$

Similar to muon decay, except that S is massive (and scalars, not fermions):

$$\Gamma_{\eta^\pm} = \frac{G_F^2}{30\pi^3} (M_{\eta^\pm} - M_S)^5$$

$$\Delta^\pm M \equiv M_{\eta^\pm} - M_S$$



but cross section would be small


# Cross sections

$$pp \rightarrow \eta^+ \eta^- + X$$

$M_s=75$  [GeV]

$M_{h1}=120$  [GeV]

$\sigma[\text{fb}]$



# Conclusions

...if scalars are dark matter...

- Scalar sector could be much more exciting than in the SM
- Possibly signals in Direct or Indirect detection experiments
- Possibly interesting signals at the LHC
- In the meantime, parts of parameter space will be excluded